

**Table 6.1**

Proposed performance indicators for priority interventions in the areas of tobacco, alcohol and obesity prevention

<b>TIER 1. HEALTH OUTCOMES (ALL TO BE REPORTED BY INDIGENOUS STATUS)</b>		
<b>OBESITY</b>	<b>TOBACCO</b>	<b>ALCOHOL</b>
Deaths attributable to overweight and obesity	Deaths attributable to tobacco	Deaths attributable to alcohol
Hospital separations attributable to overweight and obesity	Hospital separations attributable to tobacco	Hospital separations attributable to alcohol

<b>TIER 2. DETERMINANTS OF HEALTH (ALL TO BE REPORTED BY INDIGENOUS STATUS)</b>		
<b>OBESITY</b>	<b>TOBACCO</b>	<b>ALCOHOL</b>
Proportion of adults (18+ years) overweight or obese	Proportion of adults (18+ years) who are daily smokers	Proportion of adults (18+ yrs) at risk of long-term harm from alcohol
Proportion of children (12–17 years) overweight or obese	Proportion of children (12–17 years) who are daily smokers	Proportion of adults (18+ years) at risk of short-term harm from alcohol at least once per month
Proportion of adults (18+ years) eating sufficient daily serves of fruit and vegetables		Proportion of children (14–17 years) at risk of long-term harm from alcohol
Proportion of adults (18+ years) insufficiently physically active to obtain a health benefit		Proportion of children (14–17 years) at risk of short-term harm from alcohol at least once per month
Proportion of people walking, cycling or using public transport to travel to work or school		
Proportion of babies breastfed for six months or more		